

U.C. COOPERATIVE EXTENSION

SAMPLE COST TO ESTABLISH AND PRODUCE

CARROTS



IMPERIAL COUNTY – 2003

Prepared by:

Keith S. Mayberry
Herman Meister

Farm Advisor, U.C. Cooperative Extension, Imperial County
Agronomy Advisor, U.C. Cooperative Extension, Imperial County

For an explanation of calculations used for the study refer to the attached General Assumptions or call the author, Keith S. Mayberry, at the Imperial County Cooperative Extension office, (619)352-9474 or e-mail at ksmayberry@ucdavis.edu.

The University of California Cooperative Extension in compliance with the Civil Rights Act of 1964, Title IX of the Education Amendments of 1972, and the Rehabilitation Act of 1973 does not discriminate on the basis of race, creed, religion, color, national origins, or mental or physical handicaps in any of its programs or activities, or with respect to any of its employment practices or procedures. The University of California does not discriminate on the basis of age, ancestry, sexual orientation, marital status, citizenship, medical condition (as defined in section 12926 of the California Government Code) or because the individuals are disabled or Vietnam era veterans. Inquiries regarding this policy may be directed to the Personnel Studies and Affirmative Action Manager, Agriculture and Natural Resources, 2120 University Avenue, University of California, Berkeley, California 94720, (510) 644-4270.

University of California and the United States Department of Agriculture cooperating.

FOREWORD

We wish to thank growers, pest control advisors, chemical applicators and dealers, custom farm operators, fertilizer dealers, seed companies, contract harvesters, equipment companies, and the Imperial County Agricultural Commissioners office for providing us with the data necessary to compile this circular. Without them we could not have achieved the accuracy needed for evaluating the cost of production for the field crop industry in Imperial County.

The information presented herein allows one to get a "ballpark" idea of field crop production costs and practices in the Imperial County. They do not reflect the exact values or practices of any one grower, but are rather an average of countywide prevailing costs and practices. Exact costs incurred by individual growers depend upon many variables such as weather, land rent, seed, choice of agrichemicals, location, time of planting, etc. No exact comparison with individual grower practice is possible or intended. The budgets do reflect, however, the prevailing industry trends within the region.

Overhead usually includes secretarial and office expenses, general farm supplies, communications, utilities, farm shop, transportation, moving farm equipment, accountants, insurance, safety training, permits, etc. In most of the crop guidelines contained in this circular we used 13 % of the total of land preparation, growing costs and land rent to estimate overhead.

Since all of the inputs used to figure production costs are impossible to document in a single page, we have included extra expense in man-hours or overhead to account for such items as pipe setting, motor grader, water truck, shovel work, bird and rodent control, etc. Whenever possible we have given the costs of these operations per hour listed on the cultural operations page.

Not included in these production costs are expenses resulting from management fees, loans, providing supervision, or return on investments. The crop budgets also do not contain expenses encumbered for road and ditch maintenance, and perimeter weed control. If all the above items were taken into account, the budget may need to be increased by 7-15%.

Where applicable we have used terminology that is commonly used in the agricultural industry. These terms are compiled in a glossary at the end of the circular. We feel that an understanding of these terms will be useful to entry-level growers, bankers, students and visitors.

Herman S Meister & Keith S. Mayberry
(Principal researchers and editors)
Vegetable Crops and Agronomy Advisors

Contributors: Eric T. Natwick
Tom A. Turini
Jose L. Aguiar
Khaled M. Bali
Juan N Guerrero

**2002-2003 Field/Vegetable Prevailing Rate for Field Operations
IMPERIAL COUNTY**

**HEAVY TRACTOR WORK & LAND
PREPARATION**

<u>OPERATION</u>	<u>\$/ACRE</u>
Plow.....	30.50
Subsoil, 2 nd gear.....	39.00
Landplane	12.75
Triplane	11.25
Chisel 15".....	25.00
Wil-Rich chisel.....	16.00
Big Ox	24.00
Slip plow.....	41.00
Pull/disc borders	6.75
Make cross checks (taps).....	6.25
Break border	6.00
Disc, stubble	21.00
Disc, regular.....	12.50
Corrugate	11.00
Disc, regular with ring roller.....	13.50
List 30" beds 12-row	15.00
List 40" beds 8-row	15.00
Float.....	10.00
Disc, borders.....	7.00
Dump (scraper) borders	14.50

LIGHT TRACTOR WORK

Power mulch dry.....	25.00
Power mulch with herbicide	28.00
Shape 30" 6 row	10.75
Shape 40" 4 row	10.75
Plant 30" beds nonprecision	20.00
Plant 40" beds nonprecision	18.00
Precision plant 30" beds	22.00
Precision plant 40" beds	20.00
Mulch plant wheat	19.50
Plant alfalfa (corrugated).....	17.50
Plant bermudagrass (flat).....	13.75
Plant sudangrass.....	14.75
Cultivate 30" beds 4-row	16.00
Cultivate 40" beds 4-row	14.00
Spike 30" beds 4-row.....	13.25
Spike 40" beds 4-row.....	11.25
Spike and furrow out 30" 4-row	14.00
Spike and furrow out 40" 4-row	12.00
Furrow out 30" beds 4-row.....	13.25
Furrow out 40" beds 4-row.....	11.25
Lilliston 30" beds 6-row	13.00
Lilliston 40" beds 4-row	13.00
Lilliston 30" beds with/herbicides 6-row.....	15.00

Lilliston 40" beds with/herbicides 4 -row.....	15.00
Inject fertilizer & furrow out 30" beds 4-row	15.00
Inject fertilizer & furrow out 40" beds 4-row	13.00
Fertilize dry & furrow out 30" beds.....	17.00
Fertilize dry & furrow out 40" beds.....	15.00
Flat inject fertilizer NH ₃	15.00
Broadcast dry fertilizer	7.00
Ground spray 40" 8-row	12.00
Ground spray 30" 8-row	14.00
Chop cotton stalks.....	13.75

HARVEST COSTS Field Crops

	<u>BY UNIT</u>
Combine alfalfa seed	41.75/acre
Windrow alfalfa seed	17.50/acre
Rake bermudagrass	5.00/acre
Swath bermudagrass	13.50/acre
Swath sudangrass.....	11.25/acre
Rake sudangrass.....	5.25/acre
Swath alfalfa	8.00/acre
Rake alfalfa.....	4.50/acre
Bale (all types of hay- small bale)	0.65/bale
Haul & stack hay – small bale	0.25/bale
Bale (large bale 4X4).....	10.00/bale
Bale (large bale Jr. 3X4).....	9.00/bale
Stack & load large bale.....	6.00/bale
Dig sugar beets	2.60/clean ton
Haul sugar beets.....	2.45/clean ton
Combine wheat 15 per acre + 0.55 /cwt over 1 ton	
Haul wheat.....	5.50/ton
Combine bermudagrass seed 1 st time	40.00/acre
Combine bermudagrass seed 2 st time	25.00/acre
Haul bermudagrass seed (local).....	175/load
Haul bermudagrass seed (Yuma).....	300/load

MISCELLANEOUS OPERATIONS BY THE HOUR

Motor grader.....	48.00
Backhoe	45.00
Water truck	40.00
Wheel tractor	35.00
Scraper.....	36.00
Versatile.....	56.00
D-6.....	56.00
D-8.....	70.00
Buck ends of field.....	28.00
Pipe setting (2 men)	37.00
Laser	88.00
Work ends (disc out rotobucks).....	35.00

CARROT CULTURE 2002-2003

Annual acreage, yield, and value of fresh market carrots
Imperial County, CA (1997-2001)

Year	Acres	Yield/Acre*	Value/Acre
2001	4,534	800	\$5,000
2000	7,420	780	\$3,900
1999	6,798	834	\$7,681
1998	6,692	796	\$6,253
1997	6,560	777	\$5,359

Annual acreage, yield, and value of processing & other carrots
Imperial County, CA (1997-2001)

Year	Acres	Yield/Acre*	Value/Acre
2001	10,578	29.00	\$2,320
2000	11,130	30.00	\$2,400
1999	10,197	33.62	\$2,452
1998	9,724	32.66	\$2,301
1997	9,454	33.90	\$1,058

*50 pound master poly sacks containing cello packs basis. Many sold bulk.
Source: Imperial County Agricultural Commissioner's Reports 1997-2001

PLANTING-HARVESTING DATES Early maturing carrots are planted in early September. Later maturing carrots are planted in October and early November. Most of the carrots grown locally are shipped to Bakersfield for packing. The majority of the harvest starts in February and runs through mid-May. Carrots could be harvested December to early June if desired. Florida, Texas, Bakersfield and Arizona provide the most market competition during our shipping season.

VARIETIES Caro Pak; *Seminis*; Apache *Sunseeds*; Navajo *Sunseeds*; Indiana *Bejo*; and HM 02 *Harris Moran*. Varieties used for processor carrots (shortcuts, baby, whole, or cut and peel) includes: Primecut 59 *Sunseeds*; Sugar Snax *Sunseeds*; Tripleplay 58 *Sunseeds*; and Snackpack *Seminis*.

PLANTING INFORMATION Carrots may be planted 6 seed lines per 40 inch bed. Three rows are placed on each bed shoulder. Rows are usually 1½ inches apart. However, there are many variations of the number of lines, spacing between lines and bed width depending upon the shipper's needs and ultimate product use. A ballpark value is 38-42 plants per foot of bed for fresh market. For processor carrots, the plant population is 80-90 plants per foot of bed. Planting fewer seed produces large carrots for the shredder processing use, planting large amounts of seed is necessary for smaller diameter carrots for short cut use.

Natural and pelleted seed are both used. For natural seed, most growers use converted Planet Jr. planters utilizing a random flow distribution of seed drop. Stanhay planters with split shoes have been used for precision planting.

Seed is placed in a shallow groove and not covered. Enough sandy soil will cave into the groove during sprinkler irrigation to place the seed at the proper depth.

Carrots may be sown at a rate of 1,000,000 live seed per acre. Germination is often 80-90 percent and seed counts may be 175,000 to 400,000 seed per pound. The seeding rate per acre is calculated accordingly. Again this seeding rate is a variable factor and the ultimate control over population should be the decision of the shipper.

SOILS Carrots have been grown on many of the soil types in the low desert. However, best root development is obtained with the lighter, sandy-textured soils. Carrots should not be grown on stratified soils. If soils are too heavy (clayey), deformed and hairy roots will result. A deep orange colored root will not develop properly if the soil stays too wet.

IRRIGATION Carrots are sprinkler irrigated for stand establishment. Carrots germinate slowly; therefore, the beds must be kept moist to prevent crusting. They may take as long as 10 days to germinate in cool weather.

Sprinklers also reduce salinity, which is important, since carrots are very sensitive to salt. Sprinkler costs include rent, in-and-out labor, system maintenance and sprinkler operation.

After sprinkling, carrots are normally furrow irrigated for the remainder of the season. The number of irrigations may vary with climatic conditions, soil type, ultimate use of the carrot crop, and variety. The number of irrigations may vary from 4 to 6 per season after sprinkling.

Drip irrigation has not worked well with carrots. Excess water due to over-irrigation or producing carrots on poorly drained soils tends to increase the incidence of hairy roots.

If carrot fields are allowed to become too dry and then irrigated, there may be significant splitting of roots. Dryness tends to cause the cell walls to harden and lose elasticity. When more water is applied the carrot core expands while the outer layers do not, resulting in splits.

FERTILIZERS Previous crop history is helpful in determining early-season nitrogen fertilizer requirements. If the field has some residual nitrogen (N), there is no need to apply more N until the seedlings emerge. Carrot roots are vulnerable to forking if too much nitrogen is applied preplant. Phosphate is applied before listing at rates of 450 to 500 pounds of 0-45-0 per acre. Preplant fertilizer should be disced into the soil before listing to prevent forking.

Sidedress applications of 60 to 80 pounds of actual N are made during the growing season. Commonly used materials are dry ammonium nitrate (34-0-0), liquid ammonium nitrate (20-0-0), and UAN 32 (32-0-0). Shippers may wish to vary nitrogen recommendations to the grower based upon knowledge of the varietal performance and plant spacing.

Nitrogen deficiency in carrots is not readily apparent when viewing a field. Deficient fields might show an irregular pattern in height of the top growth, but the foliage will still be green in color. Since carrots are often grown on sandy soils, taking petiole analysis on a regular basis will help monitor the fertilizer status.

NEMATODE CONTROL Needle nematode (*Longidorus africanus*) and root knot nematode (*Meloidogyne* spp.) must be controlled by fumigation or chemicals to prevent forked carrot roots.

A common method of nematode control is to apply metam sodium at a rate of 60 gallons per acre in preplant flood irrigation prior to listing the beds. The soil needs to be pre-irrigated and the material applied in a second irrigation a week to ten days later. Carrot shippers have discovered that metam sodium used on carrots will provide beneficial effects to crops that are later planted in the same soil. This phenomenon is most likely due to reduction in nematode and soil borne diseases.

PESTS AND DISEASES Crickets, grasshoppers, striped flea beetle larvae and cutworms can be a problem when seedlings emerge. Later, aphids, whiteflies and spider mites may attack the leaves. Cutworms may attack crowns and have been a major problem in recent years.

Powdery mildew (*Erysiphe polygoni*) needs to be controlled if detected early in the growing season. Damping-off fungi (*Rhizoctonia solani* and *Pythium* sp.) are controlled by seed treatment. Cavity spot (*Pythium* sp.) and forking disorders have become serious problems in some fields. Root rots (*Pythium* spp.) and bacterial soft rot (*Erwinia* spp. and *Pseudomonas* spp.) are usually absent when carrots are grown in fields that have good drainage.

Early blight (*Cercospora carotae*) and late blight (*Alternaria dauci*) are fungal diseases found occasionally in carrots. These diseases must be controlled to prevent economic damage.

Black Crown (*Alternaria radicina*) is borne on seed and in soil. It has been detected in Imperial County. Plant clean seed, use crop rotation or deep till to bury the inoculum away from the crown.

HARVESTING The majority of the carrots are harvested by machines, however, a small acreage is hand bunched (with tops intact). Harvest machines can handle two full beds at a time. Loaded semi-trucks and trailers haul fresh dug carrots to local sheds or transport them to Bakersfield for washing, sizing, grading and packing. Growers must share in the freight to ship field-run carrots to Bakersfield.

The packout percentage of field-run carrots is variable depending upon variety, growing conditions, disease incidence, insect damage, mechanical damage during harvest and packing conditions. Packout rates of 70 to 80 percent are excellent, 60 to 70 percent is good, and below 55 to 60 percent is poor.

Carrots are marketed with tops on (bunched carrots), in 1-to 2- pound cello bags or topped loose in master poly containers. Prices vary according to container size.

Two sizes of cello carrots are packed: 'Jumbos' and 'Standards'. Standard cellos must be between 6 to 12 inches long and greater than 5/16 inch in diameter. There must be no less than 7 and no more than 13 carrots for a 1-pound cello. Carrots over 1½ inch in diameter at the crown are classified as Jumbos. Normally a master container with 48 "one pound" bags weighs 55 to 59 pounds as additional weight per bag.

The short-cut or baby whole carrot market has exploded in recent years. Substantial acreage is planted specifically for the processor market.

Precut carrot sticks are now being marketed nationally. Shredded carrots, matchstick cuts (Julienne cut), crinkle cut and coins are also marketed.

Bunched carrots are undercut and hand-sorted with 24 bunches per carton. The tops are bound with a wire twist tie. Master bunches are windrowed and loaded by hand onto special types of trucks which have moveable chain-link beds. Full loads are taken to sheds where electric motors are hooked up to rotate the chain-link beds (similar to a continuous conveyer belt) and unload the bunches into a water bath. This cushions the fall and reduces carrot shattering and cracking.

Bunches are washed and hand loaded into waxed cartons. Top-loaded crushed ice is used to cool the roots that are then taken to cold storage.

POSTHARVEST Mature, topped carrots have a very long shelf life if stored properly. At 34°F and 98 percent relative humidity, carrots may be stored for up to five months. Washing helps remove decay organisms and reduce loss. However, most of the carrots grown locally are shipped to retail markets soon after harvest.

Bunched carrots are highly perishable because of the tops. They may be stored for only two weeks under temperatures of 32°F and 98 percent relative humidity. Carrots should not be stored near ethylene sources (ripening fruits in particular) or they may develop a bitter flavor.

For more information on carrots, see “Carrot Production in California”, DANR Publication 7226 available from the Imperial County Cooperative Extension Office or for a free Internet download go to <http://anrcatalog.ucdavis.edu/specials.ihtml>

CARROT PROJECTED PRODUCTION COSTS 2002-2003

Hand labor at \$9.25per hour (\$6.75 plus SS, unemployment insurance, workman's compensation, transportation, supervision and fringe benefits.)

Yield--850 (39 Tons/Acre) 50-lb. Master poly containers

OPERATION	Cost	Materials		Hand Labor		Cost Per Acre
		Type	Cost	Hours	Dollars	
LAND PREPARATION						
Stubble disc 1x	21.00					21.00
Subsoil	39.00					39.00
Disc 2x	12.50					25.00
Triplane 1x	11.25					11.25
Border, cross check & break borders	19.00					19.00
Flood 1x		Water 2 ac/ft	32.00	2	18.50	50.50
Disc 2x	12.50					25.00
Apply metam sodium	30.00	metam sodium	135.00			165.00
Disc 1x	12.50					12.50
Triplane 1x	11.25					11.25
Fertilizer,spread	7.00	500 lb. @ 0-45-0	58.75			65.75
List 40" beds	15.00					15.00
TOTAL LAND PREPARATION						460.25
GROWING PERIOD						
Plant	22.00	Hybrid seed 550M	155.00			177.00
Sprinkler irrigate	185.00					185.00
Herbicide	12.50		5.00			17.50
Cultivate 2x	14.00					28.00
Spike 2x	11.25					22.50
Fertilize & furrow out 2x	13.00	150 lb. N @ .32	48.00			74.00
Herbicide, layby 1x	12.50	Lorox	25.00			50.00
Water-run fertilizer		50 lb. N @ .32	16.00			16.00
Irrigation 8X		Water 3 3/4 ac/ft	60.00	9	83.25	143.25
Disease control 1x	10.00	Fungicides	25.00			35.00
Insect control 2x	10.00	Insecticide	31.00			51.00
GROWING PERIOD						799.25
GROWING PERIOD & LAND PREPARATION COSTS						1259.50
Land Rent (net acres)						225.00
Cash Overhead	11 % of harvest costs and land rent					163.30
TOTAL PREHARVEST COSTS						1647.80
HARVEST COSTS						
Harvest by machine, haul to Bakersfield, cool, pack and sell		850 -50 lb. sacks @ (Packout 85% Cellos & 15 % Jumbos)		4.50 /sack (contract rate)		3825.00
TOTAL OF ALL COSTS						5472.80

PROJECTED PROFIT OR LOSS PER ACRE
Price/master poly sack (dollars)

Poly sacks per acre		Price/master poly sack (dollars)					Break-even \$/sack
		5.00	5.50	6.00	6.50	7.00	
750		-1273	-898	-523	-148	227	6.70
800		-1248	-848	-448	-48	352	6.56
850		-1223	-798	-373	52	477	6.44
900		-1198	-748	-298	152	602	6.33
950		-1173	-698	-223	252	727	6.23

* Harvest cost may vary with the shipper, the field conditions and the market.