


**U.C. COOPERATIVE EXTENSION**

**SAMPLE COST TO ESTABLISH AND PRODUCE**

***CABBAGE***



**IMPERIAL COUNTY – 2004**

**Prepared by:**

**Herman S Meister      Farm Advisor, U.C. Cooperative Extension, Imperial County**

---

For an explanation of calculations used for the study refer to the attached General Assumptions or call the author, Herman Meister, at the Imperial County Cooperative Extension office, (760)352-9474 or e-mail at [hmeister@ucdavis.edu](mailto:hmeister@ucdavis.edu).

The University of California Cooperative Extension in compliance with the Civil Rights Act of 1964, Title IX of the Education Amendments of 1972, and the Rehabilitation Act of 1973 does not discriminate on the basis of race, creed, religion, color, national origins, or mental or physical handicaps in any of its programs or activities, or with respect to any of its employment practices or procedures. The University of California does not discriminate on the basis of age, ancestry, sexual orientation, marital status, citizenship, medical condition (as defined in section 12926 of the California Government Code) or because the individuals are disabled or Vietnam era veterans. Inquiries regarding this policy may be directed to the Personnel Studies and Affirmative Action Manager, Agriculture and Natural Resources, 2120 University Avenue, University of California, Berkeley, California 94720, (510) 644-4270.

University of California and the United States Department of Agriculture cooperating.

---

## FOREWORD

We wish to thank growers, pest control advisors, chemical applicators and chemical dealers, custom farm operators, fertilizer dealers, seed companies, contract harvesters, equipment companies, and the Imperial County Agricultural Commissioner's office for providing us with the data necessary to compile this circular. Without their cooperation we could not have achieved the accuracy needed for evaluating the cost of production for the field crop industry in Imperial County.

The information presented herein allows one to get a "ballpark" idea of field crop production costs and practices in the Imperial County. Most of the information was collected through verbal communications via office visits and personal phone calls. The information does not reflect the exact values or practices of any one grower, but are rather an average of countywide prevailing costs and practices. Exact costs incurred by individual growers depend upon many variables such as weather, land rent, seed, choice of agrichemicals, location, time of planting, etc. No exact comparison with individual grower practice is possible or intended. The budgets do reflect, however, the prevailing industry trends within the region.

Overhead usually includes secretarial and office expenses, general farm supplies, communications, utilities, farm shop, transportation, moving farm equipment, accountants, insurance, safety training, permits, etc. Eleven to 13% of the total of land preparation, growing costs and land rent was used to estimate overhead. Hourly rates vary with each crop depending on the workman's compensation percentages.

Since all of the inputs used to figure production costs are impossible to document in a single page, we have included extra expense in man-hours or overhead to account for such items as pipe setting, motor grader, water truck, shovel work, bird and rodent control, etc. Whenever possible we have given the costs of these operations per hour listed on the cultural operations page. Some custom operators have indicated that they are instituting a "fuel surcharge" to reflect "spikes" in fuel cost.

Not included in these production costs are expenses resulting from management fees, loans, providing supervision, or return on investments. The crop budgets also do not contain expenses encumbered for road and ditch maintenance, and perimeter weed control. If all the above items were taken into account, the budget may need to be increased by 7-15%.

Where applicable we have used terminology that is commonly used in the agricultural industry. These terms are compiled in a glossary at the end of the circular. We feel that an understanding of these terms will be useful to entry-level growers, bankers, students and visitors.

Herman S Meister, Agronomy Advisor &  
Senior Editor

Contributors:

Eric T. Natwick  
Tom A. Turini  
Khaled M. Bali  
Juan N. Guerrero  
Keith Mayberry, Emeritus

**2004-2005 Tillage & Harvest Rates  
IMPERIAL COUNTY**

**HEAVY TRACTOR WORK & LAND  
PREPARATION**

<u>OPERATION</u>	<u>\$/ACRE</u>
Plow.....	32.00
Subsoil 2 <sup>nd</sup> gear.....	45.00
Subsoil 3 <sup>rd</sup> gear.....	38.00
Landplane.....	14.00
Triplane.....	12.00
Chisel 15".....	26.00
Wil-Rich chisel.....	17.00
Big Ox.....	25.00
Slip plow.....	43.00
Mark/disc borders.....	10.50
Make cross checks (taps).....	6.75
Break border.....	6.50
Stubble disc/with cultipack.....	22.50/24.50
Regular disc/with cultipack.....	13.00/15.00
List 30"-12 row/40" 8 row.....	16.50
Float.....	11.50
Dump (scraper) borders.....	18.25
Corrugate.....	14.00

**LIGHT TRACTOR WORK**

Power mulch dry.....	27.50
Power mulch with herbicide.....	31.00
Shape 30" 6-row / 40" 4-row.....	12.75/12.75
Plant sugar beets & cotton 30"/40".....	17.00/15.00
Plant vegetables.....	20.00
Mulch plant wheat.....	20.25
Plant alfalfa (corrugated).....	18.50
Plant alfalfa (beds).....	19.00
Plant bermudagrass.....	13.75
Plant with drill (sudangrass, wheat).....	14.75
Plant corn slope.....	17.00
Cultivate 30"/40" beds 4-row.....	16.00/14.00
Spike 30"/40" beds 4-row.....	13.00/11.00
Spike and furrow out 30"/40" 4-row.....	14.00/12.00
Furrow out 30"/40" beds 4-row.....	13.00/11.00
Lilliston 30" 6-row / 40" 4-row.....	14.00/14.00
Lilliston 30" 6 row/ 40" 4-row/ herb.....	15.50/15.50
Inj fert & fur out 30"/ 40" beds 4-row.....	16.50/14.50
Fertilize dry & fur out 30"/ 40" 4-row.....	17.00/15.00
Inject fertilizer flat.....	15.00
Broadcast dry fertilizer.....	8.00
Ground spray 30"/40" 8-row.....	12.00
Chop cotton stalks 30"/40"beds.....	16.00/14.00
List 80" melon beds.....	20.00
Plant 80" melon slope beds.....	22.00

Back fill furrow (melons).....9.5

Cultivate 80" melon slope beds.....	18.00
Center 80" melon beds.....	17.00
Re-run 80" melon beds.....	11.00
Inject fertilizer & furrow out 80" melon beds.....	18.00
Bust out 80" melon beds.....	12.00

**HARVEST COSTS-FIELD CROPS**

**BY UNIT**


Windrow alfalfa seed.....	17.50/acre
Combine alfalfa seed.....	41.00/acre
Swath bermudagrass.....	13.75/acre
Rake bermudagrass.....	5.50/acre
Swath sudangrass.....	11.25/acre
Rake sudangrass.....	6.00/acre
Swath alfalfa.....	8.75/acre
Rake alfalfa.....	5.00/acre
Bale (all types of hay- small bale).....	0.70/bale
Haul & stack hay – small bale.....	0.27/bale
Bale (large bale 4X4).....	7.50/bale
Haul & stack big bale.....	3.50/bale
Load with hay squeeze.....	62.50 / load
Dig sugar beets.....	2.65/clean ton
Haul sugar beets.....	2.50/clean ton
Combine wheat ....16.00 per acre + 0.60 /cwt. over 1 ton	
Haul wheat.....	5.00/ton
Combine bermudagrass seed 1st time.....	42.50/acre
Combine bermudagrass seed 2nd time.....	26.50/acre
Haul bermudagrass seed (local).....	175/load
Pick Cotton 1 <sup>st</sup> /2 <sup>nd</sup> .....	.03cts/lb/35.00/acre

**MISCELLANEOUS RATES BY THE HOUR**

**\$/HR**

Motor grader.....	55.00
Backhoe.....	50.00
Water truck.....	40.00
Wheel tractor.....	35.00
Scraper.....	36.00
Versatile.....	60.00
D-6.....	56.00
D-8.....	73.00
Buck ends of field.....	35.00
Pipe setting (2 men).....	38.00
Laser level.....	90.00
Work ends (disc out rotobucks).....	40.00

# VEGETABLE CROPS PLANTING & HARVESTING CALENDAR IMPERIAL VALLEY, CALIFORNIA






**MONTH**

- planting
- planting/harvesting
- harvesting
- \* perennial

# IMPERIAL COUNTY WEATHER

Imperial Irrigation District  
81 year average (1914-1994)



**DAYS REQUIRED FOR SEEDLING EMERGENCE\* AT VARIOUS SOIL TEMPERATURES**

Vegetable	Soil Temperature (°F)								
	32	41	50	59	68	77	86	95	104
Asparagus	NG	NG	53	24	15	10	12	20	28
Beet	/	42	17	10	6	5	5	5	/
Cabbage	/	/	15	9	6	5	4	/	/
Cantaloupe	/	/	/	/	8	4	3	/	/
Carrot	NG	51	17	10	7	6	6	9	NG
Cauliflower	/	/	20	10	6	5	5	/	/
Celery	NG	41	16	12	7	NG	NG	NG	/
Cucumbers	NG	NG	NG	13	6	4	3	3	/
Eggplant	/	/	/	/	13	8	5	/	/
Lettuce	49	15	7	4	3	2	3	NG	NG
Okra	NG	NG	NG	27	17	13	7	6	7
Onion	136	31	13	7	5	4	4	13	NG
Parsley	/	/	29	17	14	13	12	/	/
Parsnip	172	57	27	19	14	15	32	NG	NG
Peppers	NG	NG	NG	25	13	8	8	9	NG
Radish	NG	29	11	6	4	4	3	/	/
Spinach	63	23	12	7	6	5	6	NG	NG
Sweet Corn	NG	NG	22	12	7	4	4	3	NG
Tomato	NG	NG	43	14	8	6	6	9	NG
Watermelon	/	NG	/	/	12	5	4	3	/

\*planting depth = 0.5 inches; NG = no germination; / = not tested; Source: Harrington, J. F. and P. A. Minges, Vegetable Seed Germination. California Agricultural Extension Mimeo Leaflet (1954).

## SEED CALCULATIONS (M)

Number of seed (x1000) required<sup>1</sup> per acre for common plant spacing combinations within rows and between beds. Commonly coded as “M” or 1000 seed

Plant spacing within rows <sup>2</sup> (inches)	Spacing between beds <sup>3</sup> (inches)					
	30	40	42	60	66	80
<b>1</b>	209.1	156.8	149.4	104.5	95.0	78.4
<b>1.5</b>	139.4	104.5	99.6	69.7	63.4	52.3
<b>2</b>	104.5	78.4	74.7	52.3	47.5	39.2
<b>2.5</b>	83.6	62.7	59.7	41.8	38.0	31.4
<b>3</b>	69.7	52.3	49.8	34.8	31.7	26.1
<b>4</b>	52.3	39.2	37.3	26.1	23.8	19.6
<b>6</b>	34.8	26.1	24.9	17.4	15.8	13.1
<b>8</b>	26.1	19.6	18.7	13.1	11.9	9.8
<b>10</b>	20.9	15.7	14.9	10.5	9.5	7.8
<b>12</b>	17.4	13.1	12.4	8.7	7.9	6.5
<b>14</b>	14.9	11.2	10.7	7.5	6.8	5.6
<b>24</b>	8.7	6.5	6.2	4.4	4.0	3.3
<b>36</b>	5.8	4.4	4.1	2.9	2.6	2.2

<sup>1</sup> Seeds per acre was calculated assuming one seed per spacing combination. Factors influencing the actual amount of seed needed are seed delivery method and seed viability; <sup>2</sup> Values are based on beds with a single row. For multiple rows, multiply by the number of rows per bed; <sup>3</sup> Beds are measured from center to center.

Linear feet per acre for common bed widths

Bed width (inches)	Linear feet per acre
30	17,424
40	13,068
42	12,446
60	8,712
66	7,920
80	6,534

**IMPERIAL COUNTY CABBAGE PROJECTED PRODUCTION COSTS 2004-2005**

**40 Acre Field**

Hand labor at \$9.95 per hour (\$6.75 plus SS, unemployment insurance, workman's compensation and fringe benefits)

Yield--600 cartons per acre      50 lbs. per carton.      Hybrid variety

OPERATION	Cost	Materials		Hand Labor		Cost Per acre
		Type	Cost	Hours	Dollars	
<b>LAND PREPARATION</b>						
Stubble disc / ring roller	24.50					24.50
Subsoil 2nd gear	45.00					45.00
Disc 2x	13.00					26.00
Triplane	12.00					12.00
Border, cross check & break borders	23.75					23.75
Flood		Water 1 ac/ft.	16.00	1	9.95	25.95
Disc / ring roller	15.00					15.00
Triplane	12.00					12.00
Fertilize, spread	8.00	500 lb. 11-52-0	75.00			83.00
List 40" beds	16.50					16.50
<b>TOTAL LAND PREPARATION</b>						<b>283.70</b>
<b>GROWING PERIOD</b>						
Power mulch	27.50					27.50
Precision plant, & inject insecticide	20.00	Seed 104M Admire	275.00			295.00
			60.00			60.00
Weed Control- pre-emergence	12.50	Herbicide	100.00			112.50
Sprinkler irrigate	165.00					165.00
Chemigation		Insecticide	5.50			5.50
Thin				10	99.50	99.50
Cultivate 1x	14.00					14.00
Spike 2x	11.00					22.00
Fertilize & furrow out 2x	14.50	120 lb. N as UAN 32	45.60			74.60
Water-run fertilizer		60 lb. N as UAN 32	22.80			22.80
Hand weed 2x				12	119.40	119.40
Irrigate 8x		Water 3.5 ac/ft.	56.00	5	49.75	105.75
Gated pipe (harvest)	20.00					20.00
Insect control 6x	10.25	Insecticides	125.00			186.50
Disease control 1x	11.50	Fungicides	20.00			31.50
Ring roller cleanup	7.50					7.50
<b>TOTAL GROWING PERIOD</b>						<b>1,369.05</b>
<b>GROWING PERIOD &amp; LAND PREPARATION COSTS</b>						<b>1,652.75</b>
Land Rent (net acres)						225.00
Cash Overhead-----		13 % of preharvest costs & land rent				244.11
<b>TOTAL PREHARVEST COSTS</b>						<b>2,121.86</b>
<b>HARVEST COST*</b>						
Cut, pack, haul, cool and sell		600 cartons @	4.00	per carton		2,400.00
<b>TOTAL ALL COSTS</b>						<b>4,521.86</b>

**PROJECTED PROFIT OR LOSS PER ACRE**  
Price/ 50 lb. carton (dollars)

Cartons per acre		Price/ 50 lb. carton (dollars)					Break-even \$/carton
		5.00	6.00	7.00	8.00	9.00	
500		-1622	-1122	-622	-122	378	8.24
600		-1522	-922	-322	278	878	7.54
700		-1422	-722	-22	678	1378	7.03
800		-1322	-522	278	1078	1878	6.65
900		-1222	-322	578	1478	2378	6.36

\* Harvest cost varies with the shipper, the field conditions and the market



## IMPERIAL COUNTY CABBAGE CULTURE 2004-2005

Annual acreage, yield, and value of fresh market cabbage in  
Imperial County, CA (1999-2003)

Year	Acres	Yield/Acre*	Value/Acre
2003	1,245	593	\$3,066
2002	981	650	\$3,354
2001	880	710	\$4,508
2000	908	689	\$4,369
1999	1,389	522	\$2,155

\*50 lb carton equivalent

Source: Imperial County Agricultural Commissioner's Reports 1999-2003

**PLANTING-HARVESTING DATES:** Cabbage is planted from mid-September through October. Harvesting begins in December and continues through February. The average yield varies according to market price, but may reach 1,000 cartons per acre. Normally, yields reflect market demand rather than actual yield. If prices are too low, much of the harvestable crop remains in the field. Oversupply from competing markets depresses price rapidly to the point where harvesting is impractical. A shortage in the cabbage supply, however, can create windfall profits. Cabbage is a very high-risk crop!

Cabbage is often grown under contract with fast food outlets and coleslaw manufacturers. The demand for local fresh market cabbage depends upon the availability of the cabbage harvested in coastal California and Texas.

**VARIETIES:** Headstart *Seminis*; Grenadier *Syngenta*; Fast Vantage *Sakata*; Supreme Vantage *Sakata*; and Charmant *Sakata* are popular green varieties.

Commonly used red varieties include: Primero *Bejo*; Red Jewel *Sakata*; Sombrero *Bejo*; Cardinal *Harris Moran* and Red Rookie *Sakata*

**PLANTING INFORMATION:** Double seed lines on 40-inch beds are used for cabbage production. The seed is normally planted with a precision planter at 2 to 3 inches down the row spacing, at a depth of ¼ inch or less. Seed lines are usually 13 inches apart.

When seed are placed 3 inches apart within rows, on 2 seed lines per 40-inch bed, roughly 104,000 seed are required per acre. When plants develop 2 to 3 true leaves, seedlings are thinned to 12 to 14 inches within rows.



**IRRIGATION:** Sprinkler irrigation is used to germinate the crop. Once the seedlings have emerged, the field is then converted to furrow irrigation. Cabbage grows well on loam to silty clay soil. Cabbage has intermediate salt tolerance.

**FERTILIZERS:** Five hundred pounds of 11-52-0 broadcast prior to listing is standard practice. Sidedress applications of nitrogen (60-80 lb N/acre) are common. Ammonium nitrate or UAN32 solutions are often used for sidedress application.

**PESTS AND DISEASES:** Insect pests of cabbage include silverleaf whitefly, crickets, cutworms, beet armyworms, flea beetles, saltmarsh caterpillars, aphids, thrips, and cabbage looper. Once an insect burrows into cabbage heads, chemical control is nearly impossible. Neonicotinoid insecticides applied at planting are used to control silverleaf whiteflies.

Cabbage is an alternate host of the sugar beet cyst nematode (*Heterodera schachtii*) and should not be incorporated into a rotation program with sugar beets. Downy mildew (*Peronospora parasitica*) may require control if moist, cool conditions are present.

Black rot (*Xanthomonas campestris* pv. *campestris*) occurs occasionally in Imperial County. Plant using disease-free seed or disease-free transplants.

Oedema is a physiological disorder of cabbage that is manifested by pits and craters on the epidermis of the leaves. It is thought to be caused by excessive irrigation, especially during cloudy, humid weather. A similar condition may occur on the outer leaves of cabbage as a result of sand blasting.

Blind plants are created by mechanical damage or genetic defects. The damage caused by birds and insect feeding injures the growing point and a head does not develop.

**WEED CONTROL:** Herbicides are fairly effective at controlling weeds in cabbage with the exception of London rocket and shepherd's purse. Hand weeding is often necessary to remove weeds that develop during the winter.

**HARVESTING:** Fields are harvested by hand. Cabbage is packed 24 heads per bulge-packed carton. A carton may weigh over 55 pounds. Head counts can vary from 18 to 24 heads per carton, however many sales are made on the basis of net weight. Cabbage is normally sold at retail stores by the pound.

Cone-shaped cabbage heads are not acceptable in markets on the West Coast and in the Pacific Northwest. Some cabbage is grown under contract with fast food outlets for coleslaw and salad mixes.

Cabbage may be either hydrovaced or vacuum cooled. It should be stored under refrigeration after cooling.



**POSTHARVEST:** Cabbage should be stored at 32°F and a 98 percent relative humidity. Storing cabbage at low humidity causes wilting and senescence. Cabbage is sensitive to ethylene and should not be stored near ethylene sources (i.e. ripening fruits), because loss of green color and abscission of the leaves will result.

---

For more information see “Cabbage Production in California”, DANR Publication 7208 available from the Imperial County Cooperative Extension Office or for a free download from the Internet go to <http://anrcatalog.ucdavis.edu/specials.ihtml>

-----*Notes*-----



## GLOSSARY

**Air spray** The application of chemicals by aircraft.

**Back fill furrows** To shave soil off the top of melon beds and place it into a furrow in order to bring the irrigation water closer to the melon seedline.

**Bed** Mounded soil that is shaped and used for planting; beds are separated by furrows.

**Bell** Bell pepper.

**Big Ox** A chisel with 7 shanks used to rip soil 18-24 inches deep.

**Blacken the beds** To wet/darken a bed with irrigation water.

**Black Ice** Ice formation on asparagus that is clear and therefore difficult to detect.

**Blanks** Lack of individual kernel formation in corn.

**Brassicac**s Plants belonging to the genus *Brassica*, of the mustard family (Cruciferae), including cabbage, kale, broccoli, cauliflower, turnip, and mustard; all brassicas are crucifers, but not all crucifers are brassicas.

**Break a field** To harvest a crop the first time in a season.

**Break borders** To tear down flat flood borders or flat crop borders.

**Breaker** A tomato fruit that is beginning to show color change from green to pink on the blossom end; preceded by the *mature green* stage.

**Brix** A measure of sugar content, especially in tomatoes; a graduated scale, used on a hydrometer, that indicates the weight of sugar per volume of solution.

**Brown bead** A physiological disorder of broccoli thought to be related to lack of calcium uptake and excessive heat during head formation.

**Buck ends of field** The remaking of beds at the end of a field in order to channel irrigation water properly; a necessary practice when beds at the end of a field are destroyed due to insufficient turn around space for farm equipment.

**Cateye** A condition in broccoli where some beads begin breaking into yellow flower; also called *starring*.

**Cello** Poly bags which hold one or two pounds of carrots; from "cellophane".

**Chisel** A tractor-mounted, knife-like implement used to rip soil about 20 inches deep.

**'choke** Artichoke

**Cole crops** Any of various plants of the genus *Brassica*, of the mustard family.

**Cos** Romaine Lettuce

**Cross checks** Small dikes at perpendicular angles to borders used for water diversion into a field.

**Crucifers** Plants belonging to the Cruciferae or mustard family (e.g., broccoli, brussel sprouts, cabbage, cauliflower, etc.).

**Cucurbits** Plants belonging to the melon or gourd family (e.g., cantaloupe, watermelon, pumpkin, cucumbers, squash, etc.).

**Cull** To separate unwanted product from desirable product.

**Cultipacker** A farm implement used to break up clods of soil; consists of groups of knobbed metal rings stacked together.

**Cultivate** To work beds after planting in order to control weeds, loosen soil, and allow for application of fertilizer.

**Curd** The edible portion of marketed cauliflower.

**Custom rate** The value assigned to a cultural operation by farmers for cost accounting; normally includes the cost of the operator.

**Damping-off** A fungal disease of seedlings that causes rotting of the stem at the soil level and collapse of the plant.

**Doubles** The placement of two seeds rather than one when one is intended.

**Drift** Agrichemicals, dust or pests, which inadvertently fall on nearby (usually adjacent) non-target crops; usually the result of spraying products (especially products of small particle size) on windy days or of poor equipment operation.

**Drip Irrigation** The slow application of low pressure water in tubes or pipes (buried or on the surface): sometimes called trickle irrigation.

**Edema** (oedema) A physiological disorder of plant resulting from over-watering; numerous small bumps on the lower side of leaves or on stems divide, expand, and break out of the normal leaf surface and at first form greenish-white swellings or galls; the exposed surface

later becomes rusty colored and has a corky texture; especially common in cabbage.

**Excelsior** Fine wood shavings; used for stuffing, packing, etc.

**Feathering** Premature flowering of asparagus due to high temperatures.

**Flats** Flattened asparagus spears caused by certain varietal characteristics.

**Float** A large, wooden frame pulled with a tractor for rough leveling of the soil surface.

**Flood irrigation** A method of irrigation where water is applied to a field by gravity; the water is applied to a field by gravity; the water is channeled by earth borders that are usually 70 feet apart.

**'flower** Cauliflower

**Forking** The division of a tap root (especially carrots and lettuce) into branches; can be caused by nematode feeding, soil-borne pathogens, and soil texture.

**Frost kissed** Produce that has been frozen in the field and has a frosty appearance.

**Furrow irrigation** A method of irrigation where water is applied to fields by gravity flow down furrows; the water enters the bed by capillary action.

**Furrow out** The removal of soil from furrows by tractor-mounted shovels.

**Gated pipe** Large diameter pipes used to deliver low pressure water to each furrow; used to keep head end of field dry for cultivation or harvesting.

**Green line** A term used to describe the appearance of an emerging row crop as plants germinate and emerge above the soil line, a *green line* appears; often growers switch from sprinkler to furrow irrigation when a field can be *green-lined*.

**Ground spray** The application of an agrichemical by a tractor-mounted sprayer.

**Hollow stem** A physiological disorder in broccoli resulting from excessive plant spacing.

**Honeydew** Sweet excrement from aphids and whiteflies as a result of feeding on plant sap. Honeydew attracts ants and will support the growth of fungi (sooty mold).

**Hydrocool** To cool produce using ice cold water.

**Inject fertilizer** The application of liquid fertilizer in the top or sides of a bed.

**Jelly** Gelatinous material present in *mature-green* tomatoes (see also *locule*).

**Landplane** A large, tractor-pulled, land leveling machine.

**Laser level** A land surface leveler that uses a laser guiding device to maintain an accurate grade.

**Layby** To apply an herbicide or other agrichemical at the last opportunity to enter a field with a tractor prior to harvest.

**Lilliston** A rolling cultivator with curved tines which uses ground speed to assist in working up the soil surface in order to destroy weeds.

**Listing** Throwing soil in to a mound to make beds.

**Locules** Tomato fruit seed cavity.

**Mature-green** A stage of tomato fruit development when the fruit is fully grown and shows brownish ring at the stem scar after removal of the calyx; color at the blossom end has changed from light green to yellow-green and the seeds are surrounded by *jelly*.

**Motor grader** A large grader normally used to cut tail ditches for draining off excess surface water.

**Naked pack** Head lettuce packed without a wrapper.

**Pegging** the emergence of a *radicle* from seed and its placement in the soil.

**Pipe setting** Installing 2-inch plastic tubes through a soil berm with a hydraulic ram; the pipes are used to control the flow or irrigation water.

**Power mulch** A tractor-mounted, power rototiller.

**Precision planter** Planters which drop seeds at exact intervals; may function mechanically or by vacuum.

**Primed seed** Lettuce seed that has been *primed* for germination by soaking in *osmotic* solutions (e.g., polyethylene glycol [PEG]) as a preventative to *thermodormancy*.

**Pull borders** To make flood berms used to channel the water.

**Punching pipe** see *pipe setting*.

**Putting the crop to sleep** A phrase used to describe the over-watering of tomatoes by furrow irrigation following sprinkler irrigation; encourages shallow rooting and decreased plant growth.

**Radicle** The embryonic root.

**Random flow planter** A non-precision planter; seed drop is regulated by agitating the seed in a hopper over a hole; planting rate depends upon hole size and tractor speed.

**Ricing** Undesirable granulation of floret tips in cauliflower.

**Roll beds** A large, metal roller used to firm beds prior to thinning.

**Rototill** To mechanically mix soil.

**Row** A line of plants or a bed with a single line of plants.

**Seedline** A line down a bed in which seeds are planted.

**Sidedress** To place agrichemicals in a band next to a row of plants.

**Silking** Period of corn ear formation when silky threads emerge from the ear tip.

**Slant bed** A culturing technique where beds are slanted towards the winter sun (35-37 degrees from horizontal) such that the bed is perpendicular to the sun's rays.

**Slip plow** An implement pulled by a caterpillar and used to make deep cuts into the soil whereby soil from below is carried upward into the cut; used to improve drainage.

**Slush-ice-cooling** A cooling method used on broccoli; a mixture of water and ice is forced rapidly into cartons to cool the product.

**Spike** The running of tractor-mounted shanks into the soil or beds to improve aeration and drainage.

**Sprinkler irrigate** The application of irrigation water by pressurized injection into the air.

**Starring** see *cateye*

**Stinger** A root emerging from seed; a *radicle*

**Stubble disc** An implement used to chop crop residue and incorporate it into the soil; the blades are scalloped and operate like a pizza cutter.

**Subbing** Irrigation method where water is applied to a field in furrows and allowed to travel across beds by capillary action.

**Subsoil** The pulling of large, hard-faced shanks through the soil up to 42 inches deep; used to shatter soil layers and improve drainage.

**Swamper** Watermelon harvesting crew member.

**Swath** To cut a tall crop such as asparagus fern.

**Taps** See *cross checks*

**Tasseling** The emergence of corn inflorescence.

**Thermodormancy** A condition of lettuce seed where high temperatures (>86°F) make seed go dormant, thus inhibiting germination.

**Thin** The removal of excess crop plants and weeds in the seedline in order to achieve desired plant spacing.

**Tillering** Emergence of multiple stalks from the same root in corn.

**Tip burn** A condition, especially in lettuce, where leaf tips are burned; thought to be due to lack of calcium uptake; foliar applications of calcium do not correct the problem.

**Trió** A head lettuce having crew unit consisting of two cutters and a packer; only used in *naked pack* lettuce.

**Triplane** A smaller, three-wheeled version of a *landplane*.

**Triwall cardboard** Triple-layered, corrugated cardboard used to make very sturdy fiberboard containers for watermelon.

**Vacuum cooling** A cooling method whereby commodities are placed in a strong-walled room, air pressure is reduced and heat consumed in the process cools the product.

**Versatile** A large caterpillar-sized tractor with rubber tread; used to pull discs and other implements; safe for crossing asphalt roads.

**Water run** An application of an agrichemical in irrigation water (i.e., furrow irrigation).

**White star** White markings at the blossom end of tomatoes that turn from green to white as the fruit matures; an indicator of maturity in tomatoes.

**Wil-rich chisel plow** An implement used to work wet or moist soils prior to making beds.

**Wind whip** Girdling of seedling stems due to high winds. Seedlings are especially susceptible following thinning or weeding; cole crops are most susceptible.