


safflower
sample costs
and
production



Agricultural Extension Service
University of California
Imperial County
Court House, El Centro

Cost Data Sheet No. 6
UC Cooperative Extension

SAFFLOWER---SAMPLE PRODUCTION COSTS

Mechanical operations at custom rates. Hand labor at \$1.90 per hour (\$1.50 plus Social Security, unemployment insurance, and fringe benefits).

Based on 2 ton yield per acre.

OPERATION	Custom Rate	MATERIALS		HAND LABOR		SAMPLE COSTS Per Acre
		Type	Cost	Hours	Dollars	
LAND PREPARATION						
Plow	8.00					8.00
Disc 2x	2.00					4.00
Weed Control	3.00	herbicide	6.00			9.00
Float 2x	1.00					2.00
Fertilize & List	3.00	200# N (NH ₃)	12.00			15.00
TOTAL LAND PREPARATION						\$ 38.00
GROWING PERIOD						
Plant	3.00	15# seed	1.80			4.80
Irrigate 12x		water 6 ac.ft.	12.00	6.3	12.00	24.00
TOTAL GROWING PERIOD						28.80
HARVEST COSTS						
Combine	6.50/acre + .125 cwt over 1 ton					10.00
Haul	2.00 per ton					4.00
TOTAL HARVEST COSTS						14.00
Land Rent						50.00
Cash overhead 15% of above						19.62
TOTAL COSTS						\$ 150.42

Cost per Ton = \$ 75.21

SOIL REQUIREMENT

Safflower is adapted to all soils in Imperial Valley. It is moderately salt tolerant; however, for best results it should not be grown in very saline fields. The important factor is that the soil be well drained, not only on the surface but internally. Safflower will not tolerate standing water due to its susceptibility to phytophthora root rot.

LAND PREPARATION

Fields must be level enough to prevent standing water. The crop can be planted on beds, either in a mulch or in dry beds and irrigated up. It can also be planted flat in borders either in a mulch or dry bed irrigated up.

PLANTING

Optimum planting dates range from December 15th to January 15th. When planting on double row 40" beds, 15 lbs. of seed will be adequate. When safflower is drilled in flat 30 lbs. of seed should be used. Seven or 14" drill rows should be used.

When safflower is planted in a mulch, the seed should be placed $1\frac{1}{2}$ "-2" deep. A 1" planting depth is recommended for dry plantings. Safflower planted in a mulch emerges several days earlier than when planted dry and irrigated up.

VARIETIES

The variety Gila is recommended for the Imperial Valley. Frio is a new variety which shows promise of good yields accompanied by higher oil content.

IRRIGATION

For maximum yield, safflower will require 10 - 12 irrigations depending on the season and soil type. It should never be allowed to become stressed for moisture since moisture stress tends to increase the amount of root rot.

FERTILIZATION

Safflower requires 200 lbs. of nitrogen per acre for maximum yield.

Safflower has not shown a response in yield to added phosphorous fertilization.

DISEASES

Two major diseases of safflower: phytophthora root rot and safflower rust. The varieties Gila and Frio are resistant to phytophthora root rot. One method of avoiding safflower rust is to plant in fields which do not have an immediate safflower history.

INSECTS

Insect pests which may occur on safflower are: cutworm, green peach aphid, thrips, lygus bugs, loopers, stink bugs, bollworms, leafminers, and leafhoppers. Consult the Imperial County Pest Control Recommendations and University of California Agricultural Extension Service Circular 532.

HARVESTING

Safflower is ready to harvest when the leaves and particularly the bracts around the heads turn brown.

MARKETING

Most safflower is grown under contract with a processing plant for a guaranteed price.

Circular 532 by Knowles and Miller contains more comprehensive information about the crop. UC Cooperative Extension

Prepared by
Imperial County
Agricultural Extension Service
Staff
Revised November 1966

Co-operative Extension work in
Agriculture and Home Economics
College of Agriculture, University
of California, and United States
Department of Agriculture co-oper-
ating. Distributed in furtherance
of the Acts of Congress of May 8,
and June 30, 1914. George B.
Alcorn, Director, California Agri-
cultural Extension Service
UC Cooperative Extension